

LET US (S)PRAY

WHAT?	WHY?	HOW?	WHEN?		3D SPRAYING	REMARKS
			1ST SPRAYING	2D SPRAYING		
Apple	Scab	Bordeaux Mixture	Just before Blossoms Open	Just after Blossoms Drop	10 days after 2d Spraying	1st and 2d Spraying same as 2d and 3d for scab; merely add arsenate of lead to Bordeaux Do not use Lime-sulphur on growing plants
	Codling Moth	Arsenate of Lead combined with Bordeaux	Just after Blossoms Drop	10 days later	Last week of July or 1st week of August for 2d brood	
	Oyster Shell Scale	Lime-Sulphur	March or early April but before growth starts			
Cherry and Plum	Mildew and Shot-hole fungus	Bordeaux Mixture, 3-4-50	When leaves are about 1/3 grown	10 to 12 days later	10 to 12 days later	
Currant and Gooseberry	Mildew, blight and Currant worm	Bordeaux and Arsenate of Lead	When leaves are fully developed	2 to 3 weeks later		
Grapes	Mildew and Anthracnose	Bordeaux	Before leaf buds open	2 to 3 weeks later	3rd, 4th and 5th applications at intervals of 2 weeks if required	
Strawberry	Leaf-spot or blight and leaf eating insects	Bordeaux and Arsenate of Lead	When first leaves appear	After Blossoms fall		
Raspberry and Blackberry	Anthraco-nose and fungous diseases	Bordeaux	As above	2 weeks later		Spray new growth after fruit harvest